arXiv:math/9805099v1 [math.AG] 22 May 1998

DESINGULARIZED MODULI SPACES OF SHEAVES ON A $K3$, II.

May 20 1998

Kieran G. O'Grady

0. Introduction.

This is a sequel to [O4]. In that paper we constructed a symplectic desingularization of $\mathcal{M}_4$, the moduli space of rank-two semistable torsion-free sheaves F on a $K3$ surface X with $c_1(F)=0$ and $c_2(F)=4$. In this paper we will prove that our desingularization, denoted $\widetilde{\mathcal{M}}_4$, is a new (ten-dimensional) irreducible symplectic variety. Explicitely, we will show that $\widetilde{\mathcal{M}}_4$ is one-connected and that $h^{2,0}(\widetilde{\mathcal{M}}_4)=1$; this means that $\widetilde{\mathcal{M}}_4$ is an irreducible symplectic variety. Furthermore we will prove that $b_2(\widetilde{\mathcal{M}}_4)\geq 24$. Since all known ten-dimensional irreducible symplectic varieties have $b_2=7$ or $b_2=23$ [B,H], our results show that all deformations of $\widetilde{\mathcal{M}}_4$ are new irreducible symplectic varieties. The paper is organized as follows. In the first (and longest) section we prove that $\widetilde{\mathcal{M}}_4$ is connected and that $h^{2,0}(\widetilde{\mathcal{M}}_4)=1$: we adapt to our situation Jun Li's strategy for determining the stable b_1, b_2 of moduli spaces of sheaves on a surface [Li2]. In the second section we prove $\widetilde{\mathcal{M}}_4$ is simply-connected: first we prove that if $\mathcal{O}_X(1)$ has degree 2 (we can assume this by a deformation argument), $\widetilde{\mathcal{M}}_4$ is birational to a certain Jacobian fibration over $|\mathcal{O}_X(2)|$, then we show by a monodromoy argument that the Jacobian fibration is simply-connected. In the third section we exhibit a 24-dimensional subspace of $H^2(\widetilde{\mathcal{M}}_4;\mathbf{Q})$.

1. $\widetilde{\mathcal{M}}_4$ is irreducible and $h^{2,0}(\widetilde{\mathcal{M}}_4)=1$.

The proof goes as follows. First, by well-known arguments involving deformations of polarized $K3$'s [GH,O1 §2] we can assume the polarization has genus two, i.e. $H^2=2$; this will be assumed throughout the paper. We will also assume that X is a very general genus two $K3$, i.e. that $Pic(X)=\mathbf{Z}[H]$; thus H is c-generic for any c [O4,(0.2)], so that by [O4, §2] there is a symplectic desingularization $\widetilde{\pi}\colon\widetilde{\mathcal{M}}_4\to\mathcal{M}_4$. We consider the morphism $\phi\colon\mathcal{M}_4\to\mathbf{P}^N$ associated to a high power of the determinant line-bundle [LP1,Li1], and we let $\widetilde{\phi}\colon\widetilde{\mathcal{M}}_4\to\mathbf{P}^N$ be the composition with the desingularization map $\widetilde{\pi}$. We will show that $\widetilde{\phi}$ is semi-small, hence we can apply the Lefschetz Hyperplane Section (LHS) Theorem [GM] as if $\widetilde{\phi}$ were an embedding. Let $C\in|H|$ be a smooth curve: following Jun Li we choose a particular linear subspace $\Lambda\subset\mathbf{P}^N$ of codimension at most 4, such that

$$\widetilde{\phi}^{-1}\Lambda=\widetilde{V}_C\cup\widetilde{\Sigma}_C\cup\widetilde{B}_C, \tag{1.0.1}$$

where $\widetilde{V}_C,\widetilde{\Sigma}_C,\widetilde{B}_C$ are closed subsets such that:

(1) $\widetilde{V}_C\setminus(\widetilde{\Sigma}_C\cup\widetilde{B}_C)$ consists of the points x such that $\widetilde{\pi}(x)$ parametrizes a locally-free sheaf whose restriction to C is not semistable.

(2) $\widetilde{\Sigma}_C=\widetilde{\pi}^{-1}\Sigma_C$, where

$$\Sigma_C:=\{[I_Z\oplus I_W]|\ Z\text{ or }W\text{ intersects }C\}.$$

(3) $x\in\widetilde{B}_C\setminus\widetilde{\Sigma}_C$ if and only if $\widetilde{\pi}(x)$ parametrizes a stable sheaf which is not locally-free at some point of C.

This is where the hypothesis that C has genus two is used: since the moduli space of rank-two semistable vector-bundles on C with trivial determinant has dimension three, the intersection of four generic theta-divisors is empty, and this gives Decomposition (1.0.1) for a Λ of codimension at most four. By the LHS Theorem, the map induced by inclusion

$$H^q(\widetilde{\mathcal{M}}_4;\mathbf{Z})\to H^q(\widetilde{\phi}^{-1}\Lambda;\mathbf{Z})$$

is an isomorphism for $q\leq 5$. Thus we are reduced to analyzing $\widetilde{V}_C$, $\widetilde{\Sigma}_C$, $\widetilde{B}_C$, and how they intersect. The component $\widetilde{V}_C$ will be described in terms of $\mathcal{M}(H,3)$, the moduli space of semistable rank-two torsion-free sheaves E on X with $c_1(E) = H$, $c_2(E) = 3$. In fact, if $x \in \widetilde{V}_C$ is generic, and F is the sheaf parametrized by $\widetilde{\pi}(x)$, let G be the elementary modification of F associated to the desemistabilizing sequence for $F|_C$, and let $E := G(1)$; then E is a stable rank-two vector-bundle with $c_1(E) = H$, $c_2(E) = 3$, hence $[E] \in \mathcal{M}(H, 3)$. Conversely F can be reconstructed from E and the choice of a certain rank-one subsheaf of $E|_C$. Equivalently, $\widetilde{V}_C$ is birational to a moduli space of parabolic sheaves $F_0 = F \supset F_1 = G \supset F_2 = F(-C)$. Building on our knowledge of $\mathcal{M}(H, 3)$, we will show that $\widetilde{V}_C$ is birational to $\mathrm{Pic}^1(C) \times \mathcal{M}(H, 3) \times \mathbf{P}^1$. On the other hand $\widetilde{\Sigma}_C$, $\widetilde{B}_C$ are birational to $\mathbf{P}^1$-fibrations over $C \times X \times X^{(2)}$ and $C \times X^{(3)}$, respectively. We will also give birational descriptions of components of the intersection between $\widetilde{\Sigma}_C$ and $\widetilde{V}_C$, $\widetilde{B}_C$. At this point we will be ready to prove the main results of the section. First, since

$$\widetilde{V}_C \cap \widetilde{\Sigma}_C \neq \emptyset \neq \widetilde{\Sigma}_C \cap \widetilde{B}_C,$$

we get $b_0(\widetilde{\mathcal{M}_4}) = 1$. Next, if V'_C, Σ'_C, B'_C are desingularizations of $\widetilde{V}_C$, $\widetilde{\Sigma}_C$, $\widetilde{B}_C$ respectively, the pull-back map

$$\alpha: \Gamma(\Omega^2_{\widetilde{\mathcal{M}_4}}) \to \Gamma(\Omega^2_{V'_C}) \oplus \Gamma(\Omega^2_{\Sigma'_C}) \oplus \Gamma(\Omega^2_{B'_C})$$

is injective by the LHS Theorem. Of course the components of a point in $\mathrm{Im}\alpha$ must satisfy some compatibily conditions: these conditions will imply that $\mathrm{Im}\alpha$ is one-dimensional, hence $h^{2,0}(\widetilde{\mathcal{M}_4}) = 1$.

Remark. The procedure outlined above should allow us to determine the Betti numbers of $\widetilde{\mathcal{M}_4}$ up to b_5 included. However, even the determination of b_2 (which should be equal to 24) requires a much more detailed analysis of $\widetilde{V}_C \cup \widetilde{\Sigma}_C \cup \widetilde{B}_C$; the calculation of $h^{2,0}$ is simpler because $h^{2,0}$ is a birational invariant.

1.1. The determinant map ϕ.

Le Potier and Jun Li [LP1,Li1] have constructed a determinant line bundle $\mathcal{L}$ on $\mathcal{M}_4$ (in general on any moduli space of semistable torsion-free sheaves on a surface). They proved that if $m \gg 0$ then $\mathcal{L}^{\otimes m}$ is base-point free [Li1, Thm.3] and that the *determinant map*

$$\phi_m: \mathcal{M}_4 \to \mathbf{P}\left(H^0(\mathcal{M}_4, \mathcal{L}^{\otimes m})^*\right)$$

can be identified with the map to the Uhlenbeck compactification [Li1 Thm.4,FM]. From now on we fix a very large m and we set $\phi = \phi_m$. We will give an explicit description of ϕ. Let $\mathcal{M}_4^{st}, \mathcal{M}_4^{lf} \subset \mathcal{M}_4$ be the (open) subsets parametrizing stable and locally-free sheaves, respectively. By Lemma (1.1.5) of [O4],

$$\mathcal{M}_4 \setminus \mathcal{M}_4^{st} = \Sigma := \{[I_Z \oplus I_W] \mid \ Z, W \text{ zero-dimensional subschemes of } X \text{ with } \ell(Z) = \ell(W) = 2\},$$

hence $\mathcal{M}_4^{lf} \subset \mathcal{M}_4^{st}$. In particular [Li1, Thm.4] the restriction of ϕ to $\mathcal{M}^{lf}$ is an isomorphism onto its image. Let the *boundary* of $\mathcal{M}_4$ be $B := \mathcal{M}_4 \setminus \mathcal{M}_4^{lf}$, and set $B^{st} := B \cap \mathcal{M}_4^{st}$.

(1.1.1) Proposition. *Keeping notation as above, let $[F] \in B^{st}$. Then $F^{**} \cong \mathcal{O}_X^{(2)}$.*

Proof. Since F is not locally-free, $c_2(F^{**}) < c_2(F) = 4$. Thus Hirzebruch-Riemann-Roch gives $\chi(F^{**}) > 0$. By Serre duality $h^2(F^{**}) = h^0(F^*) = h^0(F^{**})$, hence $h^0(F^{**}) > 0$. A non-zero section of F^{**} must have isolated zeroes by slope-semistability, hence we have an exact sequence

$$0 \to \mathcal{O}_X \xrightarrow{\alpha} F^{**} \to I_Z \to 0,$$

where $\ell(Z) = c_2(F^{**})$. If $\ell(Z) = 0$ then $F^{**} \cong \mathcal{O}_X^{(2)}$, and we are done. By Cayley-Bacharach $\ell(Z) \neq 1$. Now assume $\ell(Z) \geq 2$: then we can find a subsheaf $I_W \subset \mathcal{O}_X$, with $\ell(W) = (4 - \ell(Z))$, such that $\alpha(I_W) \subset F$, and this contradicts stability of F. **q.e.d.**

In order to describe ϕ on the boundary we introduce the *singularity cycle* of a point $[F] \in \mathcal{M}_4$: we assume that the representative F has been chosen so that it is isomorphic to the direct sum of the successive quotients of its Harder-Narasimhan filtration: this means that if F is strictly semistable, then $F \cong I_Z \oplus I_W$. With this hypothesis we set

$$sing[F] := \sum_{p \in X} \ell_F(p),$$

where $\ell_F(p)$ is the length at p of the Artinian sheaf F^{**}/F. It follows from Lemma (1.1.5) of [O4] and Proposition (1.1.1) that if $[F] \in B$, then $sing[F] \in X^{(4)}$. By [Li1, Thm.4] the restriction of ϕ to B is simply the singularity map

$$\begin{array}{ccc} B & \xrightarrow{\phi} & X^{(4)} \\ [F] & \mapsto & sing[F] \end{array}.$$

As is easily checked the image $\phi(B)$ is equal to the whole $X^{(4)}$, hence the Uhlenbeck stratification [FM] becomes

$$\mathrm{Im}\phi = \mathcal{M}_4^{lf} \coprod X^{(4)}. \tag{1.1.2}$$

1.2. The map $\widetilde{\phi}$.

Let $\widetilde{\phi} := \phi \circ \widetilde{\pi}$, where $\widetilde{\pi} \colon \widetilde{\mathcal{M}}_4 \to \mathcal{M}_4$ is the desingularization of [O4, §2]. Let

$$\Gamma_d := \{x \in \mathrm{Im}\widetilde{\phi} | \ \dim \widetilde{\phi}^{-1}(x) = d\}.$$

(1.2.1) Proposition. *Keep notation as above. The map $\widetilde{\phi}$ is semi-small, i.e.*

$$\mathrm{cod}(\Gamma_d, \mathrm{Im}\widetilde{\phi}) \geq 2d$$

for all d.

Roughly speaking the reason is that the non-degenerate two-form $\widetilde{\omega}_4$ is in a "stratified" sense the pull-back of a two-form on $\mathrm{Im}\widetilde{\phi}$ (strictly speaking this is nonsense because $\mathrm{Im}\widetilde{\phi}$ is singular). More precisely: there is a stratification

$$X^{(4)} = \coprod_p X_p^{(4)} \tag{1.2.2}$$

indicized by partitions of 4. If $p = (p_1, p_2, p_3, p_4)$ is a partition, i.e. $\sum_{i=1}^{4} ip_i = 4$, the stratum $X_p^{(4)}$ parametrizes cycles

$$\gamma = \sum_{j=1}^{p_1} q_j^1 + 2\sum_{j=1}^{p_2} q_j^2 + \cdots,$$

where the q_j^i are pairwise distinct. Letting $X_\delta^{(s)} := \left(X^{(s)} \setminus diagonals\right)$, we have an open inclusion

$$X_p^{(4)} \subset X_\delta^{(p_1)} \times X_\delta^{(p_2)} \times \cdots,$$

in particular $X_p^{(4)}$ is smooth. Let $\omega^{(s)} \in \Gamma(\Omega^2_{X_\delta^{(s)}})$ be the holomorphic two-form obtained from ω by symmetrization, and for a partition p, let $\omega_p^{(4)}$ be the two-form on $X_p^{(4)}$ given by

$$\omega_p^{(4)} := \sum_{i=1}^{4} i\pi_i^* \omega^{(p_i)},$$

where $\pi_i: X_p^{(4)} \to X_\delta^{(p_i)}$ is the i-th projection. Stratification (1.2.2) gives a stratification of $\mathrm{Im}\widetilde{\phi} = \mathrm{Im}\phi$, in which $\mathcal{M}_4^{lf}$ is the open stratum, and the $X_p^{(4)}$ are the remaining strata. The stratified pull-back formula is the following:

$$\widetilde{\omega}_4|_{\widetilde{\phi}^{-1}X_p^{(4)}} = -\frac{1}{4\pi^2}\widetilde{\phi}^*\omega_p^{(4)}. \tag{1.2.3}$$

It is an immediate consequence of [O2,(2-9)].

Proof of Proposition (1.2.1). Equivalently, we must show that

$$\mathrm{cod}(\widetilde{\phi}^{-1}\Gamma_d, \widetilde{\mathcal{M}}_4) \geq d. \tag{1.2.4}$$

Let $V \subset \widetilde{\phi}^{-1}\Gamma_d$ be an irreducible component. If $V \cap \mathcal{M}_4^{lf} \neq \emptyset$, then $d = 0$ and there is nothing to prove. So assume $V \subset B = \widetilde{\phi}^{-1}X^{(4)}$. Then $V \cap \widetilde{\phi}^{-1}X_p^{(4)}$ is dense in V for some partition p. Let $y \in V$ be such that V is smooth at y and the restriction $\widetilde{\phi}: V \to \widetilde{\phi}(V)$ is submersive at y; it follows from (1.2.3) that

$$\widetilde{\omega}_4\left(T_yV, T_y\widetilde{\phi}^{-1}(\widetilde{\phi}(y))\right) \equiv 0.$$

Thus $\widetilde{\omega}_4$ induces a map

$$T_y\widetilde{\mathcal{M}}_4/T_yV \to T_y\widetilde{\phi}^{-1}(\widetilde{\phi}(y))^*,$$

which is surjective because $\widetilde{\omega}_4$ is non-degenerate. Hence

$$\mathrm{cod}(V, \widetilde{\mathcal{M}}_4) = \dim\left(T_y\widetilde{\mathcal{M}}_4/T_yV\right) \geq \dim T_y\widetilde{\phi}^{-1}(\widetilde{\phi}(y)) = d.$$

This proves (1.2.4). **q.e.d.**

Corollary. *Let $p = (p_1, \ldots, p_4)$ be a partition of 4. If $\gamma \in X_p^{(4)}$, then*

$$\dim\widetilde{\phi}^{-1}(\gamma) \leq 5 - \sum_{i=1}^{4} p_i. \tag{1.2.5}$$

Proof. The map

$$\widetilde{\phi}^{-1}X_p^{(4)} \xrightarrow{\widetilde{\phi}} X_p^{(4)}$$

is a locally trivial fibration in the analytic topology, hence the fibers have a constant dimension, say d. Thus $X_p^{(4)} \subset \Gamma_d$, so that

$$\mathrm{cod}(\Gamma_d, \mathrm{Im}\widetilde{\phi}) \leq \mathrm{cod}(X_p^{(4)}, \mathrm{Im}\widetilde{\phi}).$$

By Proposition (1.2.1) we get

$$2\dim\widetilde{\phi}^{-1}(\gamma) = 2d \leq \mathrm{cod}(X_p^{(4)}, \mathrm{Im}\widetilde{\phi}).$$

Writing out the codimension on the right, one gets Inequality (1.2.5). **q.e.d.**

Let $h^0(\mathcal{M}_4, \mathcal{L}^{\otimes m}) = N + 1$, so that $\widetilde{\phi}: \widetilde{\mathcal{M}}_4 \to \mathbf{P}^N$.

(1.2.6) Corollary. *Let $\Lambda \subset \mathbf{P}^N$ be a linear subspace of codimension at most c. The map*

$$H^q(\widetilde{\mathcal{M}}_4; \mathbf{Z}) \to H^q(\widetilde{\phi}^{-1}\Lambda; \mathbf{Z})$$

induced by inclusion is an isomorphism for all $q \leq (9 - c)$.

Proof. An immediate consequence of Proposition (1.2.1) and the (generalized) LHS Theorem [GM, p.150]. **q.e.d.**

1.3. Choice of Λ.

We fix once and for all a smooth curve $C \in |H|$. Let θ be line bundle on C of degree one (i.e. half the canonical degree); thus for $[F] \in \mathcal{M}_4$

$$\chi(F|_C \otimes \theta) = 0. \tag{1.3.1}$$

There is a canonical section [FM,Li1] σ_θ of the determinant line-bundle $\mathcal{L}$ such that

$$\mathit{supp}(\sigma_\theta) = \{[F] \in \mathcal{M}_4 |\ h^0(F|_C \otimes \theta) > 0\}.$$

By (1.3.1) the right-hand side has codimension at most one in every component of $\mathcal{M}_4$, but a priori it might contain a whole component. In any case there exists a linear subspace $\Lambda_\theta \subset \mathbf{P}^N$ of codimension at most one such that $\phi^{-1}\Lambda_\theta = \mathit{supp}(\sigma_\theta)$. The moduli space of rank-two semistable vector-bundles on C with trivial determinant has dimension three, hence a theorem of Raynaud [R] shows that if $\theta_1, \ldots, \theta_4$ are generic line-bundles of degree one, there is no semistable vector-bundle V with trivial determinant such that

$$h^0(V \otimes \theta_i) > 0, \qquad i = 1, \ldots, 4.$$

On the other hand, if V is a degree zero rank-two sheaf on C which is either singular or locally-free non-semistable, then $h^0(V \otimes \theta) > 0$ for any choice of a degree-one line-bundle θ on C. So let $\theta_1, \ldots, \theta_4$ be generic, and set $\Lambda := \Lambda_{\theta_1} \cap \cdots \cap \Lambda_{\theta_4}$; then

$$\phi^{-1}\Lambda = \{[F] \in \mathcal{M}_4 |\ F|_C \text{ is singular or locally-free non-semistable}\}. \tag{1.3.2}$$

Let $\Sigma_C, B_C \subset \mathcal{M}_4$ be defined as

$$\Sigma_C := \{[F] \in \Sigma |\ \mathit{sing}[F] \cap C \neq \emptyset\} \quad B_C := \{[F] \in B |\ \mathit{sing}[F] \cap C \neq \emptyset\},$$

where $\Sigma \subset \mathcal{M}_4$ is the locus parametrizing strictly semistable (i.e. non stable) sheaves. (Thus $\Sigma_C \subset B_C$.) Let $V_C^0 \subset \mathcal{M}_4$ be given by

$$V_C^0 := \{[F] \in \mathcal{M}_4^{lf} |\ F|_C \text{ is not semistable}\},$$

and let V_C be the closure of V_C^0. By Proposition (1.1.1) we can rewrite (1.3.2) as

$$\phi^{-1}\Lambda = V_C \cup B_C.$$

Let $\widetilde{V}_C, \widetilde{B}_C \subset \widetilde{\mathcal{M}}_4$ be the proper transforms of V_C and B_C respectively, and set $\widetilde{\Sigma}_C := \widetilde{\pi}^{-1}\Sigma_C$. The equality above gives

$$\widetilde{\phi}^{-1}\Lambda = \widetilde{V}_C \cup \widetilde{\Sigma}_C \cup \widetilde{B}_C,$$

hence by Corollary (1.2.5) we get the following result.

(1.3.3) Proposition. *The map*

$$H^q(\widetilde{\mathcal{M}}_4; \mathbf{Z}) \to H^q(\widetilde{V}_C \cup \widetilde{\Sigma}_C \cup \widetilde{B}_C; \mathbf{Z})$$

induced by inclusion is an isomorphism for $q \leq 5$.

1.4. Elementary modifications.

Let $[F] \in V_C^0$, and let

$$0 \to L \to F|_C \to L^{-1} \to 0 \tag{1.4.1}$$

be the desemistabilizing sequence, i.e. L is a line bundle of degree $d > 0$. Let $\iota: C \hookrightarrow X$ be inclusion, and let G be the elementary modification of F determined by (1.4.1), i.e. the sheaf fitting into the exact sequence

$$0 \to G \xrightarrow{\alpha} F \to \iota_* L^{-1} \to 0. \tag{1.4.2}$$

Then G is locally-free of rank two, and $c_1(G) = -H$, $c_2(G) = (4 - d)$. (First compute the Chern classes of $\iota_* L^{-1}$ by applying Grothendieck-Riemann-Roch to ι.)

(1.4.3) Lemma. *Keep notation as above. The vector-bundle G is slope-stable.*

Proof. Assume G is not slope-stable. Since $\mathrm{Pic}(X) = \mathbf{Z}[H]$, there exists an injection $\mathcal{O}_X(k) \hookrightarrow G$ with $k \geq 0$. Composing with α we get an injection $\mathcal{O}_X(k) \hookrightarrow F$, contradicting (Gieseker-Maruyama) semistability of F. **q.e.d.**

(1.4.4) Corollary. *Keeping notation as above, either $d = 1$ or $d = 2$.*

Proof. By Serre duality $h^2(G^* \otimes G) = h^0(G^* \otimes G)$. Since G is slope-stable $h^0(G^* \otimes G) = 1$, hence $\chi(G^* \otimes G) \leq 2$. Hirzebruch-Riemann-Roch gives $\chi(G^* \otimes G) = (4d - 6)$, thus $d \leq 2$. **q.e.d.**

By the above corollary we have a decomposition into locally closed subsets

$$V^0_C = V^0_C(1) \coprod V^0_C(2),$$

where

$$V^0_C(d) := \{[F] \in V^0_C | \ F|_C \to L^{-1} \to 0, \ L \text{ a degree-}d \text{ line-bundle}\}.$$

We will describe $V^0_C(d)$ in terms of $\mathcal{M}(H, 4-d)$, where $\mathcal{M}(H, 4-d)$ is the moduli space of semistable rank-two torsion-free sheaves on X with $c_1 = H$, $c_2 = (4-d)$. Let $\mathcal{M}(H, 4-d)^{lf} \subset \mathcal{M}(H, 4-d)$ be the subset parametrizing locally-free sheaves. Keeping notation as above, let $E := G(1)$; then $c_1(E) = H$, $c_2(E) = (4-d)$, so that by (1.4.3) $[E] \in \mathcal{M}(H, 4-d)^{lf}$. In order to reconstruct F from E, we notice that the bundle $E|_C$ comes with a canonical rank-one subsheaf: in fact the long exact sequence of $Tor(\cdot, \mathcal{O}_C)$ associated to (1.4.2) gives

$$0 \to L^{-1} \otimes \mathcal{O}_C(-C) \xrightarrow{\beta} G|_C \to L \to 0,$$

and if we tensor the above sequence with $\mathcal{O}_C(1)$ we obtain

$$0 \to L^{-1} \to E|_C \to L \otimes K_C \to 0. \tag{1.4.5}$$

One recovers F from E and (1.4.5) as follows. First notice that $F(-C) \hookrightarrow G$, secondly that the restriction to C of this inclusion has image equal to $\mathrm{Im}\beta$, thus we have an exact sequence

$$0 \to F(-1) \to G \to \iota_* L \to 0.$$

Tensoring with $\mathcal{O}_X(1)$ we see that F is the elementary modification of E associated to (1.4.5). The following result says that we can "invert" this construction.

(1.4.6) Lemma. *Let $d > 0$, and let $[E] \in \mathcal{M}^{lf}(H, 4-d)$. Assume $E|_C$ fits into Exact Sequence (1.4.5), where L is a line-bundle with $\deg L = d$. Let F be the elementary modification of E associated to (1.4.5), i.e. we have*

$$0 \to F \xrightarrow{\gamma} E \to \iota_*(L \otimes K_C) \to 0. \tag{1.4.7}$$

Then F is a rank-two slope-stable vector-bundle with $c_1(F) = 0$, $c_2(F) = 4$, and $[F] \in V^0_C(d)$.

Proof. The Chern classes of F are easily computed from the exact sequence defining F. Furthermore, applying the functor $Tor(\cdot, \mathcal{O}_C)$ to (1.4.7) we get Exact sequence (1.4.1), so all we have to prove is that F is slope-stable. Suppose F is not slope-stable. Since $\mathrm{Pic}(X) = \mathbf{Z}[H]$, there is an injection $\mathcal{O}_X(k) \hookrightarrow F$, with $k \geq 0$. Composing with γ we get $\mathcal{O}_X(k) \hookrightarrow E$; since E is semistable we must have $k = 0$, i.e. we have a non-zero section $\sigma \in H^0(E)$. The restriction of σ to C is a section of L^{-1}, hence zero because $\deg(L^{-1}) < 0$. Since $C \in |\mathcal{O}_X(1)|$ the section σ gives rise to an injection $\mathcal{O}_X(1) \hookrightarrow E$, contradicting semistability of E. **q.e.d.**

Let $\mathcal{E}_d$ be a tautological vector-bundle on $X \times \mathcal{M}^{lf}(H, 4-d)$ (it exists by [M2,(A.7)]). Let $\mathcal{Q}_C(d)$ be the relative quot-scheme of $\mathcal{E}_d|_{C \times \mathcal{M}^{lf}(H,4-d)}$ over $\mathcal{M}^{lf}(H, 4-d)$ parametrizing quotients $E|_C \to \xi$, where ξ is a rank-one sheaf of degree $(2+d)$, and let $\mathcal{Q}^0_C(d) \subset \mathcal{Q}_C(d)$ be the open subset parametrizing locally-free quotients. Lemmas (1.4.3)-(1.4.6) prove the following result.

(1.4.8). There is an isomorphism between $\mathcal{Q}^0_C(d)$ and $V^0_C(d)$, defined as follows: to a quotient $E|_C \to L \otimes K_C$, where $[E] \in \mathcal{M}(H, 4-d)^{lf}$ and L is a degree-d line-bundle, we associate the point $[F] \in V^0_C(d)$, where F is the locally-free sheaf fitting into the exact sequence

$$0 \to F \to E \to \iota_* L \otimes K_C \to 0.$$

1.5. $V^0_C(2)$ is nowhere dense.

We will prove the following.

(1.5.1) Proposition. *$V^0_C(1)$ is (open) dense in V^0_C.*

The proof of the proposition will be given at the end of this subsection. First we describe $\mathcal{M}(H, 2)$. Let $U := T_{\mathbf{P}^2}(-1)$; this a slope-stable rank-two vector-bundle on $\mathbf{P}^2$ with $c_1(U) = c_1(\mathcal{O}_{\mathbf{P}^2}(1))$, $c_2(U) = 1$. (That it is slope-stable follows from the fact that U has sections with an isolated zero.) Let

$$\psi \colon X \to |\mathcal{O}_X(1)|^* \cong \mathbf{P}^2$$

be the two-to-one branched cover, and set $W := \psi^* U$.

(1.5.2) Lemma. *Keeping notation as above, W is a slope-stable rank-two vector-bundle on X with $c_1(W) = H$, $c_2(W) = 2$. We have*

$$W|_C \cong \mathcal{O}_C \oplus K_C. \tag{1.5.3}$$

Furthermore $\mathcal{M}(H, 2) = [W]$.

Proof. Let's prove W is stable. Since U has sections with isolated zeroes, so does W, hence we have an exact sequence

$$0 \to \mathcal{O}_X \to W \to I_Z(1) \to 0,$$

where $Z \subset X$ is a zero-dimensional subscheme of length 2. Since $\mathrm{Pic}(X) = \mathbf{Z}[H]$, it follows that W is slope-stable. Now let $R := \psi(C)$, a line in $\mathbf{P}^2$. To prove (1.5.3) it suffices to verify that

$$U|_R \cong \mathcal{O}_R \oplus \mathcal{O}_R(1).$$

This follows immediately from the exact sequence

$$0 \to T_R \to T_{\mathbf{P}^2}|_R \to N_{R/\mathbf{P}^2} \to 0.$$

Let's prove the last statement. Since the expected dimension of $\mathcal{M}(H, 2)$ is zero, a result of Mukai [M2,(3.6)] gives that $\mathcal{M}(H, 2)$ consists of a single point, hence $\mathcal{M}(H, 2) = [W]$. **q.e.d.**

(1.5.4) Corollary. *Keeping notation as above,* $\dim V^0_C(2) = 5$.

Proof. According to (1.4.8) we have an identification $V^0_C(2) \cong \mathcal{Q}^0_C(2)$. Thus we have a morphism

$$\begin{array}{ccc} V^0_C(2) & \to & \mathrm{Pic}^2(C) \\ (W|_C \to L \otimes K_C \to 0) & \mapsto & [L] \end{array}.$$

The fiber over $[L]$ is the open subset of $\mathbf{P}H^0(L \otimes W|_C)$ corresponding to sections with no zeroes. It follows from (1.5.3) that

$$\dim \mathbf{P}H^0(L \otimes W|_C) = \begin{cases} 3 & \text{if } L \not\cong K_C, \\ 4 & \text{if } L \cong K_C. \end{cases}$$

The corollary follows immediately. **q.e.d.**

Proof of Proposition (1.5.1). By an argument similar to that proving Proposition (1.13) of [O3], one shows that every irreducible component of V^0_C has codimension at most $g(C)+1=3$. Since $\mathcal{M}_4$ is of pure dimension 10, every component of V^0_C has dimension at least 7. The proposition follows from Corollary (1.5.4).

1.6. Analysis of $\mathcal{M}(H,3)$.

We are mainly interested in the restriction to C of vector-bundles parametrized by $\mathcal{M}(H,3)^{lf}$: the key results are stated in Propositions (1.6.4)-(1.6.6). By [M1] the moduli space $\mathcal{M}(H,3)$ is smooth symplectic of dimension 4, and by [O1] it is irreducible. Our first goal is to define a morphism

$$\rho\colon \mathcal{M}(H,3)\to|\mathcal{O}_X(1)|. \tag{1.6.1}$$

(1.6.2) Lemma. *If $[E]\in\mathcal{M}(H,3)$ then $h^0(E)=2$.*

Proof. By Riemann-Roch $\chi(E)\geq 2$. By Serre duality $h^2(E)=\dim\operatorname{Hom}(E,\mathcal{O}_X)$, hence stability gives $h^2(E)=0$. Thus $h^0(E)\geq 2$. Let $\sigma\in H^0(E)$ be non-zero. We claim the quotient $Q:=E/\mathcal{O}_X\sigma$ is torsion-free. Suppose Q has torsion: then σ must vanish on a divisor, and since $\operatorname{Pic}(X)=\mathbf{Z}[H]$ this implies we have an injection $I_W(k)\hookrightarrow E$, where $k\geq 1$, and $W\subset X$ is a zero-dimensional subscheme. This contradicts semistability of E. Thus we have an exact sequence

$$0\to\mathcal{O}_X\to E\to I_Z(1)\to 0,$$

where $Z\subset X$ is a zero-dimensional subscheme of length $c_2(E)=3$. Since $H\cdot H=2$, we have $h^0(I_Z(1))\leq 1$, hence $h^0(E)\leq 2$. **q.e.d.**

In order to define ρ choose a basis $\{\sigma,\tau\}$ of $h^0(E)$. We claim $\sigma\wedge\tau\not\equiv 0$. Assume the contrary: then σ, τ generate a rank-one subsheaf of E with two linearly independent global sections, i.e. a sheaf isomorphic to $I_W(k)$, where $k\geq 1$ and $W\subset X$ is a zero-dimensional subscheme. This contradicts semistability of E. Thus

$$0\neq\sigma\wedge\tau\in H^0(\det E)=H^0(\mathcal{O}_X(1)).$$

We set

$$\rho([E]):=(\sigma\wedge\tau)\in|\mathcal{O}_X(1)|.$$

(1.6.3) Lemma. *For all $D\in|\mathcal{O}_X(1)|$, the fiber $\rho^{-1}D$ has pure dimension 2. In particular the map ρ is surjective.*

Proof. Since $\dim\mathcal{M}(H,3)=4$, every component of $\rho^{-1}D$ has dimension at least 2. It follows from [O2,(2-9)] that the symplectic form on $\mathcal{M}(H,3)$ is identically zero on $\rho^{-1}D$, hence every component of $\rho^{-1}D$ has dimension at most 2. **q.e.d.**

(1.6.4) Proposition. *Let $[E]\in\mathcal{M}(H,3)^{lf}$. Then $E|_C$ is not semistable if and only if $\rho([E])=C$. In this case*

$$E|_C\cong\xi\oplus(K_C\otimes\xi^{-1}),$$

where ξ is degree 3 line-bundle.

Proof. Assume $\rho([E])=C$. Choose a basis $\{\sigma,\tau\}$ of $H^0(E)$, and consider the exact sequence

$$0\to\mathcal{O}_X^{(2)}\xrightarrow{(\sigma,\tau)}E\to\iota_*\eta\to 0.$$

Restricting to C we get

$$0\to\xi\to E|_C\to\eta\to 0, \tag{1.6.5}$$

where ξ, η are rank-one sheaves. Thus ξ is locally-free of rank-one, and a Chern class computation gives $\deg\xi=3$. Since $C=(\sigma\wedge\tau)$ is smooth, σ and τ have no zeroes in common, and this implies η is also locally-free. Since $\det(E|_C)\cong K_C$, we have $\eta\cong\xi^{-1}\otimes K_C$. Exact Sequence (1.6.5) splits because $H^1(\xi^{\otimes 2}\otimes K_C^{-1})=0$. Now suppose $E|_C$ is not semistable, and let (1.6.5) be the desemistabilizing sequence, i.e. η is a line-bundle and $d=\deg\xi\geq 2$. Let F be the elementary modification of E defined by (1.6.5), i.e. we have

$$0\to F\to E\to\iota_*\eta\to 0.$$

Then $\operatorname{rk}F=2$, $c_1(F)=0$ and $c_2(F)=(3-d)\leq 1$. Arguing as in the proof of Lemma (1.4.3) we see that F is slope-semistable. Since there are no slope-semistable rank-two vector-bundles on X with $c_1=0$, $c_2=1$ (see the proof of (1.1.1)), we conclude that $c_2(F)=0$, hence $F\cong\mathcal{O}_X^{(2)}$. This means that $\rho([E])=C$. **q.e.d.**

(1.6.6) Proposition. *Let $[E] \in \mathcal{M}(H,3)^{lf}$. Assume $D = \rho([E])$ is not equal to C, and set $D \cdot C = p + p'$. There is an exact sequence*

$$0 \to \mathcal{O}_C(p) \to E|_C \to \mathcal{O}_C(p') \to 0, \tag{1.6.7}$$

which is split if $p \neq p'$.

Proof. Choose a basis $\{\sigma, \tau\}$ of $H^0(E)$. Since $\sigma(p)$, $\tau(p)$ are linearly dependent, there exists a non-trivial linear combination $\lambda\sigma + \mu\tau$ which is zero at p. Let

$$\epsilon := (\lambda\sigma + \mu\tau)|_C \in H^0(E|_C).$$

Since ϵ vanishes at p, it defines a non-zero map $\mathcal{O}_C(p) \to E|_C$. By Proposition (1.6.4) the bundle $E|_C$ is semistable, hence ϵ vanishes only at p and with multiplicity one. This gives Exact Sequence (1.6.7). Reversing the roles of p and p', we see that if $p \neq p'$ the sequence is split. **q.e.d.**

(1.6.8) Remark. The proof of Proposition (1.6.4) gives an isomorphism

$$\begin{array}{ccc} \rho^{-1}C & \overset{\sim}{\to} & \mathrm{Pic}^3(C) \\ [E] & \mapsto & [\xi] \end{array}.$$

Obviously this holds also if we replace C by any smooth $D \in |\mathcal{O}_X(1)|$. For a general D, the fiber $\rho^{-1}D$ is identified with the moduli space of rank-one torsion-free sheaves of degree 3 on D. The identification is given by associating to $[E]$ the subsheaf of $E|_D$ generated by $H^0(E)$.

1.7. Birational description of V_C.

Let $t \in \mathcal{Q}_C(1)$, and let

$$E|_C \xrightarrow{f} \xi \to 0$$

be the quotient represented by t: thus $[E] \in \mathcal{M}(H,3)^{lf}$, and ξ is a rank-one sheaf on C of degree 3. Since $E|_C$ is locally-free of degree 2, the kernel of f is a rank-one locally-free sheaf of degree (-1), say L^{-1}. Thus we can define a morphism

$$\begin{array}{ccc} \mathcal{Q}_C(1) & \xrightarrow{\varphi} & \mathrm{Pic}^1(C) \times \mathcal{M}(H,3)^{lf} \\ t & \mapsto & ([L],[E]) \end{array}.$$

We let φ_0 be the restriction of φ to $\mathcal{Q}_C^0(1)$. Clearly we have isomorphisms

$$\varphi^{-1}([L],[E]) \cong \mathbf{P}H^0(L \otimes E|_C), \tag{1.7.1}$$

$$\varphi_0^{-1}([L],[E]) \cong \mathbf{P}\{\sigma \in H^0(L \otimes E|_C)| \ \ \sigma \text{ has no zeroes}\}. \tag{1.7.2}$$

Let's examine φ over the open subset $\mathcal{U} \subset \mathrm{Pic}^1(C) \times \mathcal{M}(H,3)^{lf}$ consisting of couples $([L],[E])$ such that $h^0(L) = 0$ and $\rho([E])$ intersects C in exactly two points. Set

$$\begin{aligned} P_C &:= \varphi^{-1}(\mathcal{U}), \\ P_C^0 &:= \varphi_0^{-1}(\mathcal{U}). \end{aligned}$$

(1.7.3) Claim. *Both P_C and P_C^0 are fibrations over $\mathcal{U}$, the former with fibers $\mathbf{P}^1$, the latter with fibers $\mathbf{C}^*$. In particular* $\dim P_C = \dim P_C^0 = 7$.

Proof. Let $([L],[E]) \in \mathcal{U}$, and let $\rho([E]) \cap C = \{p, p'\}$. By (1.6.6)

$$E|_C \cong \mathcal{O}_C(p) \oplus \mathcal{O}_C(p'), \tag{1.7.4}$$

$$H^0(L \otimes E|_C) = H^0(L(p)) \oplus H^0(L(p')). \tag{1.7.5}$$

Since $K_C \sim p + p'$, and since $L \ncong [p']$, we see that $h^0(L(p)) = 1$. Similarly $h^0(L(p')) = 1$, hence the space of sections in (1.7.5) is two-dimensional. By (1.7.1) we conclude that the fibers of φ are isomorphic to $\mathbf{P}^1$. An easy argument identifies P_C with the projectivization of a direct image sheaf over $\mathcal{U}$ with fiber $H^0(L \otimes E|_C)$ over the point $([L],[E])$. This proves the result for φ. To finish the proof we remark that, since $h^0(L) = 0$, a section

$$\sigma = (\sigma_1, \sigma_2) \in H^0(L(p)) \oplus H^0(L(p'))$$

has a zero if and only if $\sigma_1 = 0$ or $\sigma_2 = 0$. **q.e.d.**

The following result will allow us to forget about the complement of P_C^0 in $\mathcal{Q}_C^0(1)$.

(1.7.6) Proposition. *Keeping notation as above, P_C^0 is dense in $\mathcal{Q}_C^0(1)$.*

Proof. This is a dimension count. Stratify the complement of $\mathcal{U}$ in $\mathrm{Pic}^1(C) \times \mathcal{M}(H,3)^{lf}$ according to the dimension of the fibers of φ_0 (see (1.7.2)). Using (1.6.4)-(1.6.6), one easily verifies that for each stratum $\mathcal{S}$,

$$\dim \varphi_0^{-1}(\mathcal{S}) < 7.$$

On the other hand (see the proof of (1.5.1)) every irreducible component of $V_C^0(1)$ has dimension at least 7. By (1.4.8) $\mathcal{Q}_C^0(1) \cong V_C^0(1)$, hence the above inequality shows that $\varphi_0^{-1}(\mathcal{S})$ is not dense in any component of $\mathcal{Q}_C^0(1)$. **q.e.d.**

Restricting the isomorphism $\mathcal{Q}_C^0(1) \xrightarrow{\sim} V_C^0(1)$ to P_C^0, we get a map

$$\psi_0 \colon P_C^0 \to V_C,$$

which is an isomorphism onto its image. The following result is an immediate consequence of (1.5.1) and (1.7.6).

(1.7.7) Corollary. *The image $\psi_0(P_C^0)$ is dense in V_C. In particular V_C is irreducible of dimension 7.*

1.8. Extension of ψ_0.

We will extend ψ_0 to a map

$$\psi \colon P_C \to V_C \tag{1.8.1}$$

Let $t \in P_C$ correspond to the quotient

$$E|_C \to \xi \to 0,$$

and let F be the associated elementary modification, i.e. we have

$$0 \to F \to E \to \iota_* \xi \to 0.$$

The sheaf F is torsion-free of rank two, with $c_1(F) = 0$, $c_2(F) = 4$. Since the family of quotient sheaves ξ form a flat family over P_C, the sheaves F fit together to give a shef $\mathcal{F}$ on $X \times P_C$, flat over P_C. Let

$$N_C := P_C \setminus P_C^0.$$

(1.8.2) Claim. *Let $t \in N_C$, and let*

$$F = \mathcal{F}|_{X \times \{t\}}.$$

Then F is a strictly (Gieseker-Maruyama) semistable sheaf. More precisely there is an exact sequence

$$0 \to I_Z \to F \to I_W \to 0, \tag{1.8.3}$$

where $Z, W \subset X$ are zero-dimensional subschemes of length 2, with $Z \neq W$.

Proof. Let $\varphi(t) = ([L],[E])$. By (1.6.6)

$$E|_C \cong \mathcal{O}_C(p) \oplus \mathcal{O}_C(p'),$$

where $p \neq p'$. We can assume (see the proof of (1.7.3)) that t corresponds to the quotient of the inclusion

$$L^{-1} \overset{\alpha}{\hookrightarrow} \mathcal{O}_C(p) \hookrightarrow E|_C.$$

The map α vanishes on a divisor $Z \subset C$ of degree 2. There is a global section $\tau \colon \mathcal{O}_X \to E$ which restricted to C gives the non-zero section of $\mathcal{O}_C(p)$ (see the proof of (1.6.6)), hence

$$\tau(I_Z)|_C = \mathrm{Im}\alpha.$$

Thus $\tau(I_Z)$ is a subsheaf of the elementary modification F, i.e. we have an exact sequence

$$0 \to I_Z \to F \to \eta \to 0,$$

for some rank-one sheaf η. We claim η is torsion-free. First, a local computation shows that η is locally-free at points of C. Secondly, since outside C the sheaves F and E are isomorphic, also η and $E/\mathrm{Im}\tau$ are isomorphic outside C. By slope-stability, the section τ has isolated zeroes, hence $E/\mathrm{Im}\tau$ is torsion-free, and we conclude that η is torsion-free outside C. Thus $\eta \cong I_W(k)$ for some integer k. Since $c_1(F) = 0$, we have $k = 0$, and since $c_2(F) = 4$ the length of W is 2. This proves F fits into Exact Sequence (1.8.3). Finally, since η is locally-free on C, W does not intersect C. Since $Z \subset C$, we conclude $Z \neq W$. **q.e.d.**

By the above claim the sheaves $\mathcal{F}|_{X\times\{t\}}$ are semistable for all $t \in P_C$, hence $\mathcal{F}$ induces a morphism $\psi: P_C \to \mathcal{M}_4$, which is equal to ψ_0 on P_C^0. Since $\mathrm{Im}\psi_0 \subset V_C$, and V_C is closed, we have $\mathrm{Im}\psi \subset V_C$. Thus we have defined the desired extension (1.8.1) of ψ_0. By Claim (1.8.2) we have

$$\psi: P_C \to V_C \setminus \Omega,$$

where we recall that Ω parametrizes strictly semistable sheaves of the form $I_Z \oplus I_Z$. Let's show that ψ lifts to a map

$$\widetilde{\psi}: P_C \to \widetilde{V}_C.$$

Outside Ω the desingularization map $\widetilde{\pi}$ is the blow-up of $(\Sigma \setminus \Omega)$. Since $\psi^{-1}(\Sigma) = N_C$, and since N_C is a divisor in the smooth variety P_C, hence Cartier, ψ lifts to $\widetilde{\mathcal{M}}_4$ by the universal property of blow-up. Obviously $\widetilde{\psi}(P_C)$ is contained in $\widetilde{V}_C$, and is dense in it. We choose a smooth projective completion V_C' of P_C such that $\widetilde{\psi}$ extends to a morphism

$$\Psi: V_C' \to \widetilde{V}_C.$$

Summing up, we have proved the following result.

(1.8.4) Proposition. *Keep notation as above. Then*

1. $\Psi: V_C' \to \widetilde{V}_C$ *is a birational morphism,*
2. $\Psi(P_C)$ *is dense in* $\widetilde{V}_C$, *in particular* $\widetilde{V}_C$ *is irreducible,*
3. $\Psi(N_C) \subset \widetilde{\Sigma}_C$, *in particular* $\widetilde{V}_C \cap \widetilde{\Sigma}_C \neq \emptyset$.

A key observation for the proof of $h^{2,0}(\widetilde{\mathcal{M}_4}) = 1$ is the following.

(1.8.5) Lemma. *The restriction map*

$$H^0(\Omega^2_{V_C'}) \to H^0(\Omega^2_{N_C})$$

is injective.

Proof. Let τ be a regular two-form on V_C' such that $\tau|_{N_C} \equiv 0$; V_C' is birational to $\mathbf{P}^1 \times \mathrm{Pic}^1(C) \times \mathcal{M}(H,3)$, hence there exists a regular two-form ϵ on $\mathrm{Pic}^1(C) \times \mathcal{M}(H,3)$ such that

$$\tau|_{P_C} = \varphi^*\epsilon.$$

Since the restriction of φ to N_C is a two-to-one covering of $\mathrm{Pic}^1(C) \times \mathcal{M}(H,3)^{lf}$, we get that ϵ is identically zero on $\mathrm{Pic}^1(C) \times \mathcal{M}(H,3)^{lf}$. But $\mathcal{M}(H,3)^{lf}$ is dense in $\mathcal{M}(H,3)$, hence $\epsilon \equiv 0$. Thus $\tau \equiv 0$. **q.e.d.**

1.9. Analysis of $\widetilde{B}_C$ and $\widetilde{\Sigma}_C$.

Consider the restriction of the determinant map ϕ (see (1.1)) to B:

$$\begin{array}{ccc} B & \xrightarrow{\phi|_B} & X^{(4)} \\ [F] & \mapsto & sing[F]. \end{array}$$

Let $X^{(4)}_{(4)} \subset X^{(4)}$ be the open stratum of Stratification (1.2.2), i.e. the partition is $(4) := (4,0,0,0)$. Set

$$\begin{aligned} B^0 :=& \phi^{-1} X^{(4)}_{(4)}, \\ \Sigma^0 :=& \Sigma \cap B^0. \end{aligned}$$

(1.9.1) Lemma. *The map* $\phi|_{B^0}: B^0 \to X^{(4)}_{(4)}$ *is a* $\mathbf{P}^1$*-fibration in the analytic topology. In particular* B^0 *is smooth of dimension* 9*. Furthermore* Σ^0 *is a smooth divisor in* B^0*.*

Proof. Let $\gamma \in X^{(4)}_{(4)}$, so $\gamma = q_1 + \cdots + q_4$, where the q_j are pairwise distinct. Let $Quot(\mathcal{O}^{(2)}_X, \gamma)$ be the quot-scheme parametrizing quotients

$$0 \to F \to \mathcal{O}^{(2)}_X \to \bigoplus_{j=1}^{4} \mathbf{C}_{q_j} \to 0.$$

Thus

$$Quot(\mathcal{O}^{(2)}_X, \gamma) \cong \mathbf{P}^1 \times \mathbf{P}^1 \times \mathbf{P}^1 \times \mathbf{P}^1.$$

Every sheaf parametrized by $\phi^{-1}\gamma$ is isomorphic to the kernel F of some quotient in $Quot(\mathcal{O}^{(2)}_X, \gamma)$. The group $\mathbf{P}\mathrm{Aut}(\mathcal{O}^{(2)}_X) \cong \mathrm{PGL}(2)$ acts on this quot-scheme: this is the diagonal action on $\mathbf{P}^1 \times \cdots \times \mathbf{P}^1$. As is easily verified semistability of a sheaf is equivalent to semistability of the corresponding point in $\mathbf{P}^1 \times \cdots \times \mathbf{P}^1$ with respect to the symmetric linearization. Hence

$$\phi^{-1}\gamma = \mathbf{P}^1 \times \mathbf{P}^1 \times \mathbf{P}^1 \times \mathbf{P}^1 // \mathrm{PGL}(2) \cong \mathbf{P}^1,$$

so that the fibers of $\phi|_{B^0}$ are $\mathbf{P}^1$'s. It is clear that $\phi|_{B^0}$ is locally trivial in the analytic topology. Finally, let $\gamma \in X^{(4)}_{(4)}$; since $\phi^{-1}\gamma \cap \Sigma$ consists of the three points representing strictly semistable orbits, $\Sigma \cap B^0$ is a smooth divisor in B^0. **q.e.d.**

Let

$$\begin{aligned} \mathcal{V}_C :=& \{\gamma \in X^{(4)}_{(4)} | \ \# \gamma \cap C = 1\}, \\ B^0_C :=& \phi^{-1} \mathcal{V}_C. \end{aligned}$$

Clearly $B^0_C \subset B_C$.

(1.9.2) Corollary.

1. *The map* $\phi|_{B^0_C}: B^0_C \to \mathcal{V}_C$ *is a* $\mathbf{P}^1$*-fibration, locally trivial in the analytic topolgy.*
2. B^0_C *is a smooth irreducible locally closed codimension one subset of* B^0*. In particular* $\dim B^0_C = 8$.
3. Σ^0 *and* B^0_C *intersect transversely (inside* B^0*), and the intersection is given by*

$$\Sigma^0_C := \{[I_Z \oplus I_W] | \ Z, W \text{ reduced disjoint}, \#Z \cap C = 1, W \cap C = \emptyset\}.$$

Proof. Item (1) follows from Lemma (1.9.1). Items (2)-(3) are easily verified. **q.e.d.**

Let $\widetilde{B}^0_C \subset \widetilde{\mathcal{M}}_4$ be the strict transform of B^0_C; clearly $\widetilde{B}^0_C \subset \widetilde{B}_C$.

(1.9.3) Proposition.

1. $\widetilde{B}^0_C$ *is open dense in* $\widetilde{B}_C$*.*
2. *The map* $\widetilde{B}^0_C \xrightarrow{\widetilde{\phi}} \mathcal{V}_C$ *is a* $\mathbf{P}^1$*-fibration, locally trivial in the analytic topology.*
3. $\widetilde{B}^0_C$ *is smooth irreducible of dimension* 8*.*

Proof. Let's prove Item (1). First we claim that every irreducible component of $\widetilde{B}_C$ has dimension at least 8. This is equivalent to showing that every component of B_C has dimension at least 8. Let

$$X^{(4)}_C := \{\gamma \in X^{(4)} | \ \mathrm{supp}\gamma \cap C \neq \emptyset\},$$

so that

$$(1.9.4). \qquad B_C = (\phi|_B)^{-1}\, X^{(4)}_C$$

Since $X^{(4)}$ is the Quotient of a smooth variety by a finite group, $X^{(4)}_C$ is a $\mathbf{Q}$-Cartier divisor in $X^{(4)}$; by (1.9.4) every component of B_C has codimension at most one in B, and this proves our claim. Now consider the stratification of $X^{(4)}_C$ induced by Stratification (1.2.2). If $\mathcal{S}_p \subset X^{(4)}_C$ is the stratum indicized by the partition $p = (p_1, \ldots, p_4)$, then by Corollary (1.2.5)

$$\dim \widetilde{\phi}^{-1}\mathcal{S}_p \leq \dim \mathcal{S}_p + 5 - \sum_{j=1}^{4} p_j.$$

Since $\dim \mathcal{S}_p = (2\sum_{i=1}^{4} p_i - 1)$, we get

$$\dim \widetilde{\phi}^{-1}\mathcal{S}_p \leq \sum_{i=1}^{4} p_i + 4.$$

The quantity on the right is strictly less than 8, unless $p = (4)$ (in which case it equals 8). Thus $\widetilde{\phi}^{-1}\mathcal{S}_{(4)}$ is dense in $\widetilde{B}_C$. Since $\mathcal{V}_C$ is dense in $\mathcal{S}_{(4)}$, this proves Item (1). To prove Item (2) notice that B^0_C does not intersect Ω, hence $\widetilde{B}^0_C$ is the blow-up of B^0_C at the (scheme-theoretic) intersection $B^0_C \cap \Sigma$. By Item (3) of Corollary (1.9.2) the intersection is a Cartier divisor, hence $\widetilde{B}^0_C \cong B^0_C$. Thus Item (2) follows from Item (1) of Corollary (1.9.2). Item (3) follows at once from Item (2). **q.e.d.**

Similar results hold for $\widetilde{\Sigma}_C$.

(1.9.5) Proposition. *$\widetilde{\Sigma}_C$ is irreducible of dimension 8.*

Proof. Consider Kirwan's desingularization $\widehat{\pi}\colon \widehat{\mathcal{M}}_4 \to \mathcal{M}_4$ (see [O4,(1.8)]), and let $\widehat{\Sigma}_C := \widehat{\pi}^{-1}\Sigma_C$. Let $\alpha\colon \widehat{\mathcal{M}}_4 \to \widetilde{\mathcal{M}}_4$ be the contraction along $\widehat{\Omega}$ defined in [O4,(2)]. Since $\widetilde{\Sigma}_C = \alpha(\widehat{\Sigma}_C)$, it suffices to prove $\widehat{\Sigma}_C$ is irreducible of dimension 8. Letting $\widehat{\Sigma} := \widehat{\pi}^{-1}\Sigma$, it follows from [O4,(1.7)] that $\widehat{\Sigma}$ is a $\mathbf{P}^1$-fibration over

$$\Sigma_R//\mathrm{PGL}(N) = Bl_\Omega(\Sigma).$$

Since the inverse image of Σ_C under the blow-up $Bl_\Omega(\Sigma) \to \Sigma$ is irreducible of dimension 7, we conclude that $\widehat{\Sigma}_C$ is irreducible of dimension 8. **q.e.d.**

Let $\widetilde{\Sigma}^0_C := \widetilde{\pi}^{-1}\Sigma^0_C$. The following result is an immediate consequence of Proposition (1.9.5).

(1.9.6) Corollary. *$\widetilde{\Sigma}^0_C$ is a smooth open dense subset of $\widetilde{\Sigma}_C$.*

Let $S_C := \widetilde{\Sigma}^0_C \cap \widetilde{B}^0_C$.

(1.9.7) Proposition. *S_C is a section of the $\mathbf{P}^1$-fibration $\widetilde{\Sigma}^0_C \to \Sigma^0_C$.*

Proof. This is an immediate consequence of Item (3) of Corollary (1.9.2). **q.e.d.**

Let Σ'_C be a desingularization of $\widetilde{\Sigma}_C$. Since by (1.9.6), $\widetilde{\Sigma}^0_C$ is a smooth open subset of $\widetilde{\Sigma}_C$, we can think $\widetilde{\Sigma}^0_C \subset \Sigma'_C$, hence also $S_C \subset \Sigma'_C$. The following immediate consequence of Proposition (1.9.7) will be important in the proof that $h^{2,0}(\widetilde{\mathcal{M}}_4) = 1$.

(1.9.8) Corollary. *Keeping notation as above, the restriction map*

$$H^0(\Omega^2_{\Sigma'_C}) \to H^0(\Omega^2_{S_C})$$

is injective.

1.10. Proof that $b_0(\widetilde{\mathcal{M}}_4) = h^{2,0}(\widetilde{\mathcal{M}}_4) = 1$.

Let's prove $\widetilde{\mathcal{M}}_4$ is connected. By Proposition (1.3.3) it suffices to show that

$$\widetilde{V}_C \cup \widetilde{\Sigma}_C \cup \widetilde{B}_C \tag{1.10.1}$$

is connected. Each of $\widetilde{V}_C, \widetilde{\Sigma}_C, \widetilde{B}_C$ is irreducible, by (1.8.4), (1.9.5), and (1.9.3) respectively. By Proposition (1.8.4) $\widetilde{V}_C \cap \widetilde{\Sigma}_C \neq \emptyset$, and clearly $\widetilde{\Sigma}_C \cap \widetilde{B}_C \neq \emptyset$. Thus (1.10.1) is connected. Now we prove that $h^{2,0}(\widetilde{\mathcal{M}}_4) = 1$. Let

$$\Psi \colon V'_C \to \widetilde{V}_C$$

be as in (1.8.4), and let

$$f \colon \Sigma'_C \to \widetilde{\Sigma}_C \qquad g \colon B'_C \to \widetilde{B}_C$$

be desingularizations of $\widetilde{\Sigma}_C$, $\widetilde{B}_C$ respectively. By Proposition (1.3.3) the map

$$\begin{array}{ccc} H^{2,0}(\widetilde{\mathcal{M}}_4) & \to & H^{2,0}(V'_C) \oplus H^{2,0}(\Sigma'_C) \oplus H^{2,0}(B'_C) \\ [\tau] & \mapsto & (\Psi^*[\tau], f^*[\tau], g^*[\tau]) \end{array} \tag{1.10.2}$$

is injective. We claim that also g^* is injective. So let τ be a two-form on $\widetilde{\mathcal{M}}_4$, and assume that $g^*\tau \equiv 0$. If S_C is as in (1.9.7), then

$$f^*\tau|_{S_C} \equiv g^*\tau|_{S_C} \equiv 0.$$

By Corollary (1.9.8) we conclude that $f^*\tau \equiv 0$. Letting $N_C \subset P_C$ be as in (1.8.4), it follows from $f^*\tau \equiv 0$ that $\Psi^*\tau|_{N_C} \equiv 0$. Thus Lemma (1.8.5) gives that $\Psi^*\tau \equiv 0$. By injectivity of (1.10.2) we get that $\tau \equiv 0$. This proves g^* is injective. We claim that $h^{2,0}(B'_C) = 1$. In fact, by Proposition (1.9.3) there is a rational dominant map

$$B'_C \cdots > X^{[3]}$$

with generic fiber isomorphic to $\mathbf{P}^1$, and since $h^{2,0}(X^{[3]}) = 1$ it follows that $h^{2,0}(B'_C) = 1$. Since g^* is injective we get that $h^{2,0}(\widetilde{\mathcal{M}}_4) \leq 1$. But we know $h^{2,0}(\widetilde{\mathcal{M}}_4) \geq 1$, hence $h^{2,0}(\widetilde{\mathcal{M}}_4) = 1$.

2. $\widetilde{\mathcal{M}}_4$ is simply-connected.

We will show that $\widetilde{\mathcal{M}}_4$ is birational to an open subset $\mathcal{J}$ of the Jacobian fibration parametrizing (stable) degree-six line-bundles on curves in $|\mathcal{O}_X(2)|$. This implies that $\pi_1(\mathcal{J})$ surjects onto $\pi_1(\widetilde{\mathcal{M}}_4)$, hence it will suffice to show that $\mathcal{J}$ is simply connected. The latter result is proved by a monodromy argument.

2.1. The Jacobian fibration.

Let $C \subset |\mathcal{O}_X(2)|$ be a reduced irreducible curve, $\iota \colon C \hookrightarrow X$ be inclusion, and L a degree-six line-bundle on C. We let $\mathcal{N}$ be Simpson's moduli space [LP2,S] of pure one-dimensional sheaves ξ on X, stable with respect to $\mathcal{O}_X(1)$, with

$$\chi(\xi(n)) = \chi(\iota_* L(n)).$$

Thus $[\iota_* L]$ is a typical point of $\mathcal{N}$. There is a morphism

$$\begin{array}{ccc} \mathcal{N} & \xrightarrow{\rho} & |\mathcal{O}_X(2)| \cong \mathbf{P}^5 \\ \xi & \mapsto & \mathrm{supp}\xi. \end{array} \tag{2.1.1}$$

By results of Mukai [M1] we know that $\mathcal{N}$ is smooth of dimension 10. We will be interested in a certain open subset of $\mathcal{N}$ defined as follows. First, let $U \subset |\mathcal{O}_X(2)|$ be the open subset parametrizing reduced curves. The following result is a straightforward application of the definition of stability according to Simpson [LP2,S].

(2.1.2) Lemma. *Let $[C] \in U$, and let L be a line-bundle on C such that:*

1. *if C is irreducible, the degree of L is 6,*
2. *if $C = C_1 \cup C_2$ and $L_i := L|_{C_i}$, the degree of L_i is 3.*

Then $\iota_* L$ is a stable pure one-dimensional sheaf on X.

(2.1.3) Definition. Let $\mathcal{J} \subset \mathcal{N}$ be the open set parametrizing sheaves $\iota_* L$, where $[C] \in U$, $\iota: C \hookrightarrow X$ is inclusion, and L is a line-bundle on C satisfying the hypotheses of Lemma (2.1.2). We will often denote by (C, L) the point $[\iota_* L] \in \mathcal{J}$.

Since $\mathcal{J}$ is open in $\mathcal{N}$, the results of Mukai mentioned above give the following.

(2.1.4) Proposition. *$\mathcal{J}$ is smooth of dimension* 10.

2.2. $\widetilde{\mathcal{M}}_4$ is birational to $\mathcal{J}$.

Notice that if $(C, L) \in \mathcal{J}$, then $\chi(L) = 2$, hence $h^0(L) \geq 2$. Let

$$\mathcal{J}^0 := \{(C, L) \in \mathcal{J} |\ \ C \text{ is smooth, } h^0(L) = 2 \text{ and } L \text{ is globally generated}\}.$$

As is easily verified $\mathcal{J}^0$ is open and dense in $\mathcal{J}$. Let $(C, L) \in \mathcal{J}^0$. Following Lazarsfeld [La] we will associate to (C, L) a stable rank-two vector-bundle on X with $c_1 = 0$, $c_2 = 4$; this construction will define an isomorphism between $\mathcal{J}^0$ and an open subset of $\widetilde{\mathcal{M}}_4$. Since L is globally generated, the evaluation map $H^0(L) \otimes \mathcal{O}_X \to \iota_* L$ is surjective: let E be the sheaf on X fitting into the exact sequence

$$0 \to E \xrightarrow{\epsilon} H^0(L) \otimes \mathcal{O}_X \to \iota_* L \to 0. \tag{2.2.1}$$

(2.2.2) Lemma. *Keeping notation as above, E is a slope-stable rank-two vector bundle on X with Chern classes $c_1(E) = -2H$, $c_2(F) = 6$.*

Proof. The Chern classes are easily computed from (2.2.1). To show stability, consider the exact sequence

$$0 \to L^{-1} \to H^0(L) \otimes \mathcal{O}_C \to L \to 0. \tag{2.2.3}$$

By (2.2.1) we have

$$\mathrm{Im}(\epsilon|_C) = L^{-1}. \tag{2.2.4}$$

Now suppose E is not stable: since $\mathrm{Pic}(X) = \mathbf{Z}[H]$ there is an injection of sheaves $\mathcal{O}_X(k) \overset{\alpha}{\hookrightarrow} E$, where $k \geq -1$. By (2.2.4)

$$(\epsilon \circ \alpha)|_C \in H^0(L^{-1}(-k)).$$

Since $\mathrm{deg} L = 6$, we have $\mathrm{deg} L^{-1}(-k) \leq -2$. Thus $\epsilon \circ \alpha$ vanishes on C, i.e.

$$\epsilon \circ \alpha \in H^0(\mathcal{O}_X^{(2)}(-k)(-C)) = H^0(\mathcal{O}_X^{(2)}(-k-2)).$$

Since $k \geq -1$, this last group is zero, contradiction. **q.e.d.**

Set $F := E(1)$; by the above lemma F is a slope-stable rank-two vector-bundle on X with $c_1(F) = 0$, $c_2(F) = 4$. Thus we can define $\Phi^0: \mathcal{J}^0 \to \mathcal{M}_4$ by setting

$$\begin{array}{ccc} \mathcal{J}^0 & \xrightarrow{\Phi^0} & \mathcal{M}_4 \\ (C, L) & \mapsto & [F]. \end{array}$$

We will identify an open subset $\mathcal{M}_4^0$ of $\mathcal{M}_4$ such that Φ^0 is an isomorphism onto $\mathcal{M}_4^0$. Let $[F] \in \mathcal{M}_4^{lf}$, and set $G = F(1)$. Then $c_1(G) = 2H$, $c_2(G) = 6$; by Riemann-Roch we get $\chi(G) = 2$. By Serre duality $h^2(G) = h^0(G^*)$, hence stability gives $h^2(G) = 0$. Thus $h^0(G) \geq 2$, therefore the locus where $h^0(G) = 2$ is open in $\mathcal{M}_4$. If $h^0(G) = 2$, consider the evaluation map

$$H^0(G) \otimes \mathcal{O}_X \xrightarrow{f} G.$$

Since G is slope-stable and $\mathrm{Pic}(X) = \mathbf{Z}[H]$, the determinant of f is not identically zero, so $(\mathrm{det} f) \in |\mathcal{O}_X(2)|$. We set

$$\mathcal{M}_4^0 := \{[F] \in \mathcal{M}_4^{lf} |\ \ h^0(F(1)) = 2 \text{ and } (\mathrm{det} f) \text{ is smooth}\}.$$

Clearly $\mathcal{M}_4^0$ is open in $\mathcal{M}_4$.

(2.2.5) Proposition. *The map* Φ^0 *is an isomorphism onto* $\mathcal{M}_4^0$.

Proof. Let's show $\Phi^0(\mathcal{J}^0) \subset \mathcal{M}_4^0$. Tensoring (2.2.1) by $\mathcal{O}_X(2)$ and observing that $G = E(2)$, we get

$$0 \to G \to \mathcal{O}_X(2)^{(2)} \to \iota_* L(2) \to 0. \tag{2.2.6}$$

The long exact sequence of $Tor(\cdot, \mathcal{O}_C)$ associated to (2.2.6) gives

$$0 \to L \to G|_C \to L^{-1}(2) \to 0.$$

From (2.2.6) we get a natural injection $\beta\colon \mathcal{O}_X(2)^{(2)}(-C) \hookrightarrow G$, and $\mathrm{Im}(\beta|_C) = L$. Thus we have an exact sequence

$$0 \to \mathcal{O}_X^{(2)} \xrightarrow{\gamma} G \to \iota_* L^{-1}(2) \to 0. \tag{2.2.7}$$

By adjunction $\mathcal{O}_C(2) \cong K_C$, hence Serre duality gives $h^0(L^{-1}(2)) = h^1(L)$. By hypothesis $h^0(L) = \chi(L)$, hence $h^1(L) = 0$. Thus the cohomology long exact sequence of (2.2.7) gives $h^0(G) = 2$. Furthermore the map γ drops rank along C, which is smooth, hence $[F] \in \mathcal{M}_4^0$. Now we define an inverse

$$(\Phi^0)^{-1}\colon \mathcal{M}_4^0 \to \mathcal{J}^0.$$

Let $[F] \in \mathcal{M}_4^0$, and set $G = F(1)$. Since $[F] \in \mathcal{M}_4^0$, we have an exact sequence

$$0 \to H^0(G) \otimes \mathcal{O}_X \xrightarrow{f} G \to \iota_* \xi \to 0, \tag{2.2.8}$$

where $(\det f) \in |\mathcal{O}_X(2)|$ is smooth: set $C := (\det f)$. Since C is smooth, the map f is nowhere zero, hence ξ is a line-bundle on C: we can write $\xi = L^{-1} \otimes K_C$, where L is a line-bundle. By a Chern class computation one gets $\deg L = 6$. Since $h^0(G) = 2$, the cohomology long exact sequence of (2.2.8) gives $h^0(L^{-1} \otimes K_C) = 0$. Thus $h^1(L) = 0$, so that $h^0(L) = 2$. Furthermore, the long exact sequence of $Tor(\cdot, \mathcal{O}_C)$ associated to (2.2.8) gives

$$0 \to L \to G|_C \to L^{-1} \otimes K_C \to 0.$$

Since $(\mathrm{Im} f)|_C = L$, the line-bundle L is generated by global sections. We have proved $(C, L) \in \mathcal{J}^0$. As is easily verified the map

$$\begin{array}{ccc} \mathcal{M}_4^0 & \to & \mathcal{J}^0 \\ [F] & \to & (C, L) \end{array}$$

is the inverse of Φ^0. **q.e.d.**

We can view Φ^0 as a map to $\widetilde{\mathcal{M}}_4$, because $\mathcal{M}_4^0$ is in the stable locus of $\mathcal{M}_4$. Since $\widetilde{\mathcal{M}}_4$ is irreducible, Proposition (2.2.5) implies that Φ^0 extends to a birational map

$$\Phi\colon \mathcal{J} \cdots > \widetilde{\mathcal{M}}_4.$$

Let $I(\Phi) \subset \mathcal{J}$ be the indeterminacy locus of Φ. By Proposition (2.1.4) $\mathcal{J}$ is smooth, hence $I(\Phi)$ has codimension at least two and the map induced by inclusion

$$\pi_1(\mathcal{J} \setminus I(\Phi)) \to \pi_1(\mathcal{J})$$

is an isomorphism. On the other hand, since Φ is a birational map and $\widetilde{\mathcal{M}}_4$ is smooth, the map

$$\pi_1(\mathcal{J} \setminus I(\Phi)) \to \pi_1(\widetilde{\mathcal{M}}_4)$$

is surjective. Thus we have proved the following result.

(2.2.9) Proposition. *The map Φ induces a surjection $\pi_1(\mathcal{J}) \to \pi_1(\widetilde{\mathcal{M}}_4)$.*

2.3. $\mathcal{J}$ is simply-connected.

Let $f: X \to |\mathcal{O}_X(1)|^* \cong \mathbf{P}^2$ be the two-to-one cover, and let $B \subset \mathbf{P}^2$ be the (sextic) branch curve. Let $V \subset U$ be the open subset parametrizing smooth curves $C \in |\mathcal{O}_X(2)|$. Then

$$V = U \setminus (\Delta \cap \Lambda), \tag{2.3.1}$$

where

$$\begin{aligned} \Delta :=& \{C \in |\mathcal{O}_X(2)| \mid f(C) \text{ is a smooth conic tangent to } B\}, \\ \Lambda :=& \{C \in |\mathcal{O}_X(2)| \mid f(C) \text{ is a singular reduced conic}\}. \end{aligned}$$

Let $\rho: \mathcal{J} \to U$ be the canonical map (2.1.1), and set $\mathcal{J}_V := \rho^{-1}V$. By Proposition (2.1.4) $\mathcal{J}$ is smooth, hence the map

$$j_\#: \pi_1(\mathcal{J}_V) \to \pi_1(\mathcal{J})$$

induced by inclusion $j: \mathcal{J}_V \hookrightarrow \mathcal{J}$ is surjective. We will show $j_\#$ is trivial: this will prove $\mathcal{J}$ is simply-connecetd. The map $\mathcal{J}_V \to V$ is a fibration with fibers 5-dimensional Jacobians. The homotopy exact sequence of a fibration gives an exact sequence

$$\pi_1(Jac(C)) \to \pi_1(\mathcal{J}_V) \to \pi_1(V) \to \{1\},$$

where $C \in |\mathcal{O}_X(2)|$ is a fixed smooth curve.

(2.3.2) Lemma. *Keeping notation as above, the restriction of $j_\#$ to $\pi_1(Jac(C))$ is trivial.*

Proof. We have an isomorphism $\pi_1(Jac(C)) \cong H_1(C;\mathbf{Z})$. As is easily seen the vanishing cycles on C for the family of curves parametrized by V generate all of $H_1(C;\mathbf{Z})$. Since $j_\#$ is trivial on vanishing cycles, the lemma follows. **q.e.d.**

By the above lemma, $j_\#$ induces a surjective homomorphism $\overline{j}_\#: \pi_1(V) \to \pi_1(\mathcal{J})$. We will finish the proof by showing that $\overline{j}_\#$ is trivial. This is a consequence of the following easy result.

(2.3.3) Lemma. *Let $D \subset \mathbf{C}$ be a disc centered at 0, and let $\psi: D \hookrightarrow U$ be a (holomorphic) embedding such that:*

1. *$\psi(D^0) \subset V$, where $D^0 := D \setminus \{0\}$,*
2. *$\psi(0)$ is a smooth point of Δ (or Λ), and D intersects Δ (respectively Λ) transversely. Then, after shrinking D, we can assume there exists a lift $\widetilde{\psi}: D \to \mathcal{J}$ of ψ.*

Proof. Let $q: \mathcal{C} \to D$ be the family of curves in $|\mathcal{O}_X(2)|$ parametrized by D. By Item (2) the analytic surface $\mathcal{C}$ is smooth. Shrinking D we can assume there exists a line-bundle $\mathcal{L}$ on $\mathcal{C}$ such that for each $t \in D$, the couple $(q^{-1}t, \mathcal{L}|_{q^{-1}t})$ satisfies the hypotheses of Lemma (2.1.2). By the modular property of $\mathcal{J}$ the couple $(\mathcal{C}, \mathcal{L})$ induces a morphism $\widetilde{\psi}: D \to \mathcal{J}$ lifting ψ. **q.e.d.**

To finish the proof that $\overline{j}_\#$ is trivial, let $R \subset U|$ be a straight line (notice that $|\mathcal{O}_X(2)| \setminus U$ has codimension 3) transverse to Δ and Λ. The map induced by inclusion

$$\pi_1(R \setminus (\Delta \cup \Lambda)) \to \pi_1(V)$$

is a surjection by Bertini's Theorem [GM, p.151]. The fundamental group on the left is generated by $\pi_1(D^0_{x_i})$, where $R \cap (\Delta \cup \Lambda) = \{x_1, \ldots, x_k\}$, and $D^0_{x_i} \subset (R \setminus (\Delta \cup \Lambda))$ is a small punctured disc centered at x_i. By Lemma (2.3.3) the map $\overline{j}_\#$ is trivial on each $\pi_1(D^0_{x_i})$. This shows $\overline{j}_\#$ is trivial, and concludes the proof that $\mathcal{J}$ is simply-connecetd.

3. Proof that $b_2(\widetilde{\mathcal{M}}_4) \geq 24$.

Let $\phi: \mathcal{M}_4 \to \mathbf{P}^N$ be the determinant map (1.1). Composing Donaldson's map $\mu: H_2(X;\mathbf{Q}) \to H^2(\phi(\mathcal{M}_4);\mathbf{Q})$ (see [FM,Mo]) with pull-back by ϕ, we get

$$\phi^*\mu: H_2(X;\mathbf{Q}) \to H^2(\mathcal{M}_4;\mathbf{Q}).$$

This map is an injection, because by [FM,Mo]

$$\int_{\mathcal{M}_4} \wedge^{10}(\phi^*\mu(\alpha)) = \frac{10!}{5!2^5}\left(\int_X \wedge^2\alpha\right)^5. \tag{3.1}$$

Consider the boundary divisor $B \subset \mathcal{M}_4$ (see (1.1)). By Lemma (1.9.1) B contains $\mathbf{P}^1$'s contracted by ϕ. Hence, if $h \in H^2(\mathcal{M}_4)$ is the class of a hyperplane, $\mathbf{Q}h$ and $\phi^*\mu(H_2(X;\mathbf{Q}))$ span a 23-dimensional subspace of $H^2(\mathcal{M}_4)$. Now we pull back by $\widetilde{\pi}: \widetilde{\mathcal{M}}_4 \to \mathcal{M}_4$.

(3.2) Claim. *Keep notation as above. Then*

$$\dim\left(\mathbf{Q}\widetilde{\pi}^*h + \widetilde{\pi}^*\phi^*\mu(H_2(X;\mathbf{Q}))\right) = 23.$$

Proof. The subspace $\widetilde{\pi}^*\phi^*\mu(H_2(X;\mathbf{Q}))$ has dimension 22 by (3.1). If $\widetilde{\mathbf{P}}^1 \subset \widetilde{\mathcal{M}}_4$ is the proper transform of a $\mathbf{P}^1 \subset B$ contracted by ϕ, then

$$\langle \widetilde{\pi}^*h, \widetilde{\mathbf{P}}^1\rangle \neq 0, \qquad \langle \widetilde{\pi}^*\phi^*\mu(H_2(X;\mathbf{Q})), \widetilde{\mathbf{P}}^1\rangle = 0.$$

Thus $\mathbf{Q}\widetilde{\pi}^*h$ is not contained in $\widetilde{\pi}^*\phi^*\mu(H_2(X;\mathbf{Q}))$. **q.e.d.**

Finally let's show that $\mathbf{Q}c_1(\widetilde{\Sigma})$ and $\mathbf{Q}\widetilde{\pi}^*h + \widetilde{\pi}^*\phi^*\mu(H_2(X;\mathbf{Q}))$ span a 24-dimensional subspace of $H^2(\widetilde{\mathcal{M}}_4;\mathbf{Q})$. Let $\widetilde{\Sigma}_{Z,W} \subset \widetilde{\mathcal{M}}_4$ be the fiber of $\widetilde{\pi}$ over $[I_Z \oplus I_W]$, where $Z \neq W$. By [O4, (2.3.1)] $\widetilde{\Sigma}_{Z,W} \cong \mathbf{P}^1$, and by [O4,(2.3.2)]

$$\langle c_1(\widetilde{\Sigma}), \widetilde{\Sigma}_{Z,W}\rangle = -2, \qquad \langle \mathbf{Q}\widetilde{\pi}^*h \oplus \widetilde{\pi}^*\phi^*\mu(H_2(X;\mathbf{Q})), \widetilde{\Sigma}_{Z,W}\rangle = 0.$$

Thus $\mathbf{Q}c_1(\widetilde{\Sigma})$ is not contained in $\mathbf{Q}\widetilde{\pi}^*h + \widetilde{\pi}^*\phi^*\mu(H_2(X;\mathbf{Q}))$. By Claim (3.2) we conclude that

$$\dim\Big(\mathbf{Q}c_1(\widetilde{\Sigma}) + \mathbf{Q}\widetilde{\pi}^*h + \widetilde{\pi}^*\phi^*\mu(H_2(X;\mathbf{Q}))\Big) = 24.$$

References.

[B] A. Beauville. *Variétés Kählériennes dont la première classe de Chern est nulle*, J. Differential Geom. 18 (1983), 755-782.

[FM] R. Friedman-J. Morgan. *Smooth four-manifolds and complex surfaces*, Ergeb. Math. Grenzgeb. (3. Folge) 27, Springer (1994).

[GH] L. Göttsche, D. Huybrechts. *Hodge numbers of moduli spaces of stable bundles on $K3$ surfaces*, preprint, MPI/94-80.

[GM] M. Goresky, R. MacPherson. *Stratified Morse Theory*, Ergeb. Math. Grenzgeb. (3. Folge) 14, Springer (1988).

[H] D. Huybrechts. *Compact Hyperkähler manifolds: basic results*, alg-geom/9705025.

[La] R. Lazarsfeld. *Brill-Noether-Petri without degenerations*, J. of Differential Geom. 23 (1986), 299-307.

[LP1] J. Le Potier. *Fibré déterminant et courbes de saut sur les surfaces algébriques*, Complex projective geometry, London Math. Soc. Lecture Note Series 179, Cambridge University Press (1992).

[LP2] J. Le Potier. *Systèmes cohérents et structures de niveau*, Astérisque 214, Soc. Math. de France (1993).

[Li1] J. Li. *Algebraic geometric interpretation of Donaldson's polynomial invariants of algebraic surfaces*, J. Diff. Geom. 37 (1993), 417-466.

[Li2] J. Li. *The first two Betti numbers of the moduli spaces of vector bundles on surfaces*, preprint (1995).

[M1] S. Mukai. *Symplectic structure of the moduli space of sheaves on an abelian or $K3$ surface*, Invent. math. 77 (1984), 101-116.

[M2] S. Mukai. *On the moduli space of bundles on $K3$ surfaces, I*, in Vector bundles on algebraic varieties, T.I.F.R., Oxford Univ. Press (1987), 341-413.

[O1] K. G. O'Grady. *The weight-two Hodge structure of moduli spaces of sheaves on a K3 surface*, J. of Algebraic Geom. 6 (1997), 599-644.

[O2] K. G. O'Grady. *Moduli of vector-bundles on surfaces*, Algebraic Geometry Santa Cruz 1995, Proc. Symp. Pure Math. vol. 62, Amer. Math. Soc. (1997), 101-126.

[O3] K. O'Grady. *Moduli of vector bundles on projective surfaces: some basic results*, Invent. math. 123 (1996), 141-207.

[O4] K. O'Grady. *Desingularized moduli spaces of sheaves on a $K3$, I*, preprint Dip.to di Matematica "G. Castelnuovo" 98/21 (1998).

[R] M. Raynaud. *Sections des fibrés vectoriels sur une courbe*, Bull. Soc. Math. Fr. 110 (1982), 103-125.

[S] C. Simpson. *Moduli of representations of the fundamental group of a smooth projective variety I*, Publ. Math. Inst. Hautes Études Sci. 79 (1994), 47-129.

Universitá di Roma "La Sapienza"

Dipartimento di Matematica "G. Castelnuovo"

00185 Roma

ITALIA

e-mail: ogrady@mat.uniroma1.it